



## Corrigendum: a mixed Boolean and deposit model for the modeling of metal pigments in paint layers

Enguerrand Couka, François Willot, Dominique Jeulin

### ► To cite this version:

Enguerrand Couka, François Willot, Dominique Jeulin. Corrigendum: a mixed Boolean and deposit model for the modeling of metal pigments in paint layers. *Image Analysis & Stereology*, 2015, 34 (3), pp.145. 10.5566/ias.1417 . hal-01232489

**HAL Id: hal-01232489**

**<https://hal-mines-paristech.archives-ouvertes.fr/hal-01232489>**

Submitted on 27 Nov 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CORRIGENDUM: A MIXED BOOLEAN AND DEPOSIT MODEL FOR THE MODELING OF METAL PIGMENTS IN PAINT LAYERS

ENGUERRAND COUKA, FRANÇOIS WILLOT AND DOMINIQUE JEULIN

Center for Mathematical Morphology, MINES ParisTech, PSL Research University, 35 rue Saint-Honoré, 77300 Fontainebleau, France  
e-mail: enguerrand.couka@mines-paristech.fr, francois.willot@mines-paristech.fr, dominique.jeulin@mines-paristech.fr  
(Submitted)

The expressions for the distribution function and the cumulative distribution of chord lengths given in (Couka *et al.*, 2015) (Eqs. 3 and 4) are incorrect. The latter must be replaced by:

$$P_1(\ell) = P\{L < \ell\} = 1 - \sqrt{1 - \left(\frac{\ell}{D}\right)^2}, \quad (1)$$

$$P_2(\ell) = 1 - \frac{\ell}{D} \left[ \sqrt{\left(\frac{D}{\ell}\right)^2 - 1} - \cos^{-1}\left(\frac{\ell}{D}\right) \right], \quad (2)$$

respectively. The above expressions are numerically very similar to that given in (Couka *et al.*, 2015) and to the measurements carried out on the SEM images (see updated Fig. 1 (a) and (b), which replace Figs. 7 and 18, resp).




Fig. 1. (a): Cumulative distribution of the lengths of the particles along their main direction, on a 2D section: measurements (red), theoretical distribution for cylinders with constant (green) or uniform diameter (blue). (b): Cumulative distribution function of the length of particles along the main directions as observed on SEM images (blue) and stack model (red). Theoretical formula seen in Eq. (2) shown in green.

## REFERENCES

Couka E, Willot F, Jeulin D (2015). A mixed boolean and deposit model for the modeling of metal pigments in paint layers. Image Anal Stereol 34(2):125-134.